

Topic Area: Cancer

CA-1: Reduce the death rate due to all cancers

| | |
|------------------|---|
| Target: | 161.5 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 179.4 per 100,000 standard population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-2: Reduce the death rate due to lung cancer

| | |
|------------------|---|
| Target: | 42.0 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 46.7 per 100,000 standard population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-3: Reduce the death rate due to female breast cancer

| | |
|------------------|---|
| Target: | 23.5 per 100,000 standard female population (age-adjusted) |
| Baseline: (Year) | 26.1 per 100,000 standard female population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-4: Reduce the death rate due to uterine cervix cancer

| | |
|------------------|---|
| Target: | 1.9 per 100,000 standard female population (age-adjusted) |
| Baseline: (Year) | 2.1 per 100,000 standard female population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-5: Reduce the death rate due to colorectal cancer

| | |
|------------------|--|
| Target: | 15.8 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 18.6 per 100,000 standard population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, |

| | |
|--|--|
| | Center for Health Statistics, New Jersey Department of Health |
|--|--|

CA-6: Reduce the death rate due to oropharyngeal cancer

| | |
|------------------|---|
| Target: | 2.0 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 2.2 per 100,000 standard population (age-adjusted) (2005-2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-7: Reduce the death rate due to prostate cancer

| | |
|------------------|---|
| Target: | 21.2 per 100,000 standard male population (age-adjusted) |
| Baseline: (Year) | 23.6 per 100,000 standard male population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-8: Reduce the death rate due to melanoma cancer

| | |
|------------------|---|
| Target: | 2.4 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 2.7 per 100,000 standard population (age-adjusted) (2007) |
| Data source: | Death Certificate Database, Center for Health Statistics, New Jersey Department of Health |

CA-9: Reduce the incidence rate of invasive colorectal cancer

| | |
|------------------|---|
| Target: | 39.7 per 100,000 standard population (age-adjusted) |
| Baseline: (Year) | 44.7 per 100,000 standard population (age-adjusted) (2009 preliminary) |
| Data source: | New Jersey State Cancer Registry, Cancer Epidemiology Services, New Jersey Department of Health |

CA-10: Reduce the incidence rate of invasive uterine cervical cancer

| | |
|---------|--|
| Target: | 7.2 per 100,000 standard population (age-adjusted) |
|---------|--|

| | |
|------------------|---|
| Baseline: (Year) | 8.5 per 100,000 standard female population (age-adjusted) (2009 preliminary) |
| Data source: | New Jersey State Cancer Registry, Cancer Epidemiology Services, New Jersey Department of Health |

CA-11: Reduce the incidence rate of late-stage female breast cancer

| | |
|------------------|---|
| Target: | 43.7 per 100,000 standard female population (age-adjusted) |
| Baseline: (Year) | 46.9 per 100,000 standard female population (age-adjusted) (2009 preliminary) |
| Data source: | New Jersey State Cancer Registry, Cancer Epidemiology Services, New Jersey Department of Health |

CA-12: Increase the proportion of oral and pharyngeal cancers detected at the earliest stage

| | |
|------------------|---|
| Target: | 36.9 percent |
| Baseline: (Year) | 31.5 percent (2009 preliminary) |
| Data source: | New Jersey State Cancer Registry, Cancer Epidemiology Services, New Jersey Department of Health |

CA-13: Increase the proportion of cancer survivors who are living 5 years or longer after diagnosis

| | |
|------------------|---|
| Target: | 74.6 percent |
| Baseline: (Year) | 66.6 percent (diagnosed in 2003-2007 and followed through 2010) |
| Data source: | New Jersey State Cancer Registry, Cancer Epidemiology Services, New Jersey Department of Health |

CA-14: Increase the proportion of women aged 21 to 65 years who receive a cervical cancer screening based on the most recent guidelines

| | |
|------------------|---|
| Target: | To be determined |
| Baseline: (Year) | Not yet available (2009-2011) |
| Data source: | New Jersey Behavioral Risk Factor Survey, Center for Health Statistics, New Jersey Department of Health |

CA-15: Increase the proportion of adults aged 50 to 75 years who receive a colorectal cancer screening based on the most recent guidelines

| | |
|------------------|---|
| Target: | To be determined |
| Baseline: (Year) | Not yet available (2009-2011) |
| Data source: | New Jersey Behavioral Risk Factor Survey, Center for Health Statistics, New Jersey Department of Health |

CA-16: Increase the proportion of women aged 50 to 74 years who receive a breast cancer screening based on the most recent guidelines

| | |
|------------------|---|
| Target: | To be determined |
| Baseline: (Year) | Not yet available (2009-2011) |
| Data source: | New Jersey Behavioral Risk Factor Survey, Center for Health Statistics, New Jersey Department of Health |

CA-17: Increase the proportion of men aged 40 and over whose doctor, nurse, or other health professional have ever talked to them about the advantages of the prostate specific antigen (PSA) test to screen for prostate cancer.

| | |
|------------------|---|
| Target: | To be determined |
| Baseline: (Year) | Not yet available (2009-2011) |
| Data source: | New Jersey Behavioral Risk Factor Survey, Center for Health Statistics, New Jersey Department of Health |

CA-18: Reduce the proportion of adults aged 18 years and older who report sunburn

| | |
|------------------|---|
| Target: | To be determined |
| Baseline: (Year) | Not yet available (2009-2011) |
| Data source: | New Jersey Behavioral Risk Factor Survey, Center for Health Statistics, New Jersey Department of Health |